

COVID-19 Real-time RTPCR: Does Positivity on Follow up RTPCR Always Imply Infectivity?

Tejas Suri^a MD, Saurabh Mittal^b MD DM, FCCP, Pawan Tiwari^b MD DM, Anant Mohan^c MD PhD, Vijay Hadda^d MD, Karan Madan^d MD DM, Randeep Guleria^e MD DM

^aSenior Resident, ^bAssistant Professor, ^cProfessor and Head, ^dAssociate Professor,

Department of Pulmonary, Critical Care and Sleep Medicine, All India Institute of Medical

Sciences (AIIMS), Ansari Nagar, New Delhi, India

^eProfessor, Department of Pulmonary, Critical Care and Sleep Medicine; Director, All India Institute of Medical Sciences (AIIMS), Ansari Nagar, New Delhi, India

Correspondence Address:

Dr. Pawan Tiwari,

Room No. 2, Third Floor, Porta Cabin, New Private Room Block, All India Institute of Medical

Sciences,

New Delhi – 110029, India

Email address: pavan14281@gmail.com

Telephone Number: +911126593488

Word Count: 515

Conflicts of Interest: None for all the authors

Financial Disclosures / Funding: None for all the authors

Contribution: All authors contributed equally to the manuscript.

This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives License 4.0 (<http://creativecommons.org/licenses/by-nc-nd/4.0/>). For commercial usage and reprints please contact Diane Gern (dgern@thoracic.org).

To the Editors:

We read with interest the study by De Chang et al wherein they have addressed the important question relating the duration of COVID-19 symptoms with SARS-CoV-2 viral positivity on real-time PCR. Among 16 patients with predominantly mild disease, they found that the median duration of symptoms was 8 days. They found that 8 patients remained real-time PCR positive beyond symptom resolution for a median of 2.5 days after clinical recovery (1). These findings are in congruence with the work of Young et al who reported that SARS-CoV-2 remained detectable by PCR in nasopharyngeal swabs for a median duration of 12.5 days after symptom onset. Furthermore, viral positivity by PCR outlived symptoms in half of their patients as well (2).

We noted that in the present study, De Chang et al have erroneously pointed to an instance of disease transmission from a patient after symptom resolution. However, on perusal of the cited report, contact between the index case and secondary cases occurred prior to the onset of symptoms in the index case. The discovery of the contagion occurred subsequent to the recovery of the index case who remained PCR positive on nasopharyngeal swab and sputum at this time (3). High communicability of COVID-19 in the pre-symptomatic and early symptomatic phase has been pointed out in epidemiologic modelling studies also (4).

The natural question which emerges is where does this leave us in terms of contagiousness of patients who have symptomatically recovered? While there is no definitive answer to this, many organizations have promulgated a test-based discharge strategy. This implies that patients may be discharged from hospital only after they have two negative PCR test results 24 hours apart on their nasopharyngeal swabs to prevent disease transmission. However, there is already a shortage of testing to detect new cases in most countries around the world, thus placing immense strain on a stretched public health response.

In this regard, it must be stressed that positivity on PCR does not necessarily translate to infectivity in respiratory viral illnesses. Inagaki et al have demonstrated in a ferret model, that transmission of

influenza occurred till only 5 days after the experimental infection of donor ferrets. This period correlated with culture positivity and antigen detectability. In contrast, PCR positivity continued for 11-13 days post-inoculation of the donor ferrets (5).

Interestingly, Wölfel et al examined 9 patients with mild COVID-19 infection for the correlation between PCR and viral culture results. They found that despite symptomatic recovery, the patients continued to have detectable SARS-CoV-2 RNA by PCR on nasopharyngeal swab for around 2 weeks. On the contrary, peak viral shedding occurred in early course of illness; all viral cultures beyond day 8 were negative despite positive PCR. Further, SARS-CoV-2 genomic replication was not observed beyond one week on respiratory samples (6). This coincides with the findings of He et al that patients are unlikely to be infectious beyond 1 week despite positive PCR (4). Therefore, we need to find newer tools to judge infectivity beyond symptom resolution and revise the hospital discharge (or ending isolation) criteria, especially in mild cases who have recovered clinically.

References

1. Chang D, Mo G, Yuan X, Tao Y, Peng X, Wang FS, Xie L, Sharma L, Dela Cruz CS, Qin E. Time Kinetics of Viral Clearance and Resolution of Symptoms in Novel Coronavirus Infection. [letter] *Am J Respir Crit Care Med*. 2020;201:1150-1152.
2. Young BE, Ong SWX, Kalimuddin S, Low JG, Tan SY, Loh J, et al. Epidemiologic Features and Clinical Course of Patients Infected With SARS-CoV-2 in Singapore. *JAMA*. 2020 Mar 3;
3. Rothe C, Schunk M, Sothmann P, Bretzel G, Froeschl G, Wallrauch C, et al. Transmission of 2019-nCoV Infection from an Asymptomatic Contact in Germany. *N Engl J Med*. 2020 05;382(10):970–1.
4. He X, Lau EHY, Wu P, Deng X, Wang J, Hao X, et al. Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nat Med*. 2020 Apr 15;
5. Inagaki K, Song M-S, Crumpton J-C, DeBeauchamp J, Jeevan T, Tuomanen EI, et al. Correlation Between the Interval of Influenza Virus Infectivity and Results of Diagnostic Assays in a Ferret Model. *J Infect Dis*. 2016 Feb 1;213(3):407–10.
6. Wölfel R, Corman VM, Guggemos W, Seilmaier M, Zange S, Müller MA, et al. Virological assessment of hospitalized patients with COVID-2019. *Nature*. 2020 Apr 1;